

A New Approach to the Spin $\frac{1}{2}$ Quantum Heisenberg Model

K.-H. Mütter*

*Physics Department, University of Wuppertal
D-5600 Wuppertal 1, Germany*

(WU B 93-11, Mai 93, cond-mat/9303039)

An evolution equation for the expectation values of the Boltzmann factor between monomer valence bond states is derived. It contains the whole information on the thermodynamical and magnetic properties of the spin $\frac{1}{2}$ quantum Heisenberg model. A new approximation scheme is proposed and studied in the thermodynamical limit.

I. INTRODUCTION

In 1938 Hulthén¹ determined eigenvalues and eigenstates of the spin $\frac{1}{2}$ quantum Heisenberg model on small clusters with $V \leq 10$ sites. At that time, computers were not yet at his disposal. He therefore exploited the symmetries of the model-including the conservation of total spin $\tilde{S}$ - in order to reduce the dimensions of the block Hamiltonians as much as possible. With the advent of powerful computers exact diagonalizations on larger clusters up to $V = 36$ sites² and quantum Monte Carlo simulations on much larger systems³ could be performed. Many of these computations were devoted to the determination of the ground-state properties of the antiferromagnetic model: the energy and staggered magnetization⁴, spin-spin correlators⁵ etc. Since the ground-state is known to have spin 0⁶ it is sufficient to diagonalize the Hamiltonian in this sector. Conservation of total spin, however, is not used in most of these computations. An exception is Andersons resonating valence bond (RVB) state⁷. It seems that Hulthén's construction of spin 0 states by means of valence bonds is not so suited for numerical calculations on large systems for the following reasons:

- The set of valence bond states is overcomplete; there are linear dependences among them.
- Valence bond states are not orthogonal to each other.

There have been made some attempts to overcome these deficiencies⁸. E.g. one can try to restrict the number of 'allowed' valence bond configurations in order to avoid over completeness. Such a procedure was successful for $d = 1$ ⁸. For $d > 1$, however, it looks very unpleasant: The apparent symmetries of the system -translations, rotations, reflections are hidden in complicated relations.

In this paper I want to draw the attention to some astonishing properties of the projection operators on valence bond states:

- Completeness
- Spectral like representations of the Hamiltonian and other operators.
- Anticommutation relations with the Hamiltonian.

The properties are discussed in section 2 and extended to projection operators on spin S states in section 3. In section 4 these properties are shown to describe the evolution of the expectation values of the Boltzmann factor between monomer valence bond states. In section 5 an approximation scheme for the evaluation of the evolution equation expansion is proposed: the monomer dimer cluster expansion. The thermodynamical limit of the lowest order approximation is investigated in section 6.

II. VALENCE BOND PROJECTION OPERATORS (VBPO)

We consider a lattice with an even number $V = 2v$ of sites. Spin operators $\vec{\sigma}(x)$ live at each site x . The operator

$$(x, y) = \frac{1}{4}(1 - \vec{\sigma}(x)\vec{\sigma}(y)) \quad (2.1)$$

projects the two spin state at sites x and y onto its valence bond (i.e. spin 0) part:

$$[x, y] = \frac{1}{\sqrt{2}}(\chi_+(x)\chi_-(y) - \chi_-(x)\chi_+(y)). \quad (2.2)$$

Each partition of the lattice into pairs $[x_j, y_j]$ of sites defines a valence bond configuration (VBC) K . We associate to each configuration K a state:

$$|K\rangle = [x_1, y_1][x_2, y_2]\dots[x_v, y_v], \quad (2.3)$$

with total spin 0. These valence bond states yield an overcomplete non orthogonal basis in the total spin 0 sector. In the following we are concerned with the valence bond projection operators (VBPO):

$$P(K) = |K\rangle\langle K| = (x_1, y_1)(x_2, y_2)\dots(x_v, y_v). \quad (2.4)$$

The VBPO have the following properties:

$$\text{tr } P(K) = 1, \quad (2.5)$$

$$\sum_K P(K) = a_0 1_0, \quad (2.6)$$

where 1_0 is the projection operator onto the sector with total spin 0. The ‘completeness relation’ (2.6) is proven in appendix A. If we take the trace of eq.(2.6) we get for the constant a_0 :

$$a_0 = \frac{V!!}{n_0(V)}, \quad (2.7)$$

where $V!!$ is the number of valence bond configurations and $n_0(V)$ the number of spin 0 states on a lattice with $V = 2v$ sites. In appendix A it is shown that spin-spin correlators (x, y) as they are defined in eq.(2.1) have a particular simple expansion in terms of the VBPO’s:

$$1_0(x, y)1_0 = \frac{V+2}{4a_0} \sum_K \delta([x, y] \in K) P(K). \quad (2.8)$$

According to eq.(2.8) all configurations K with a fixed valence bond $[x, y]$ contribute with equal weight. The remaining configurations have zero weight. From eq.(2.8) we obtain the expansion of the nearest neighbour Heisenberg Hamiltonian projected onto the spin 0 sector:

$$H_0 = 1_0 \sum_{\langle x, y \rangle} 4(x, y)1_0, \quad (2.9)$$

$$H_0 = \frac{V+2}{a} \sum_K N_1(K) P(K), \quad (2.10)$$

where

$$N_1(K) = \sum_{\langle x, y \rangle} \delta([x, y] \in K) \quad (2.11)$$

is the number of dimers -these are the nearest neighbour valence bonds- on the configuration K . Eq.(2.10) looks like a spectral representation. But it is not, since the valence bond states $|K\rangle$ are not orthogonal. The Hamiltonian H_0 obeys anticommutation relations with the VBPO’s which are proven in appendix A as well:

$$\frac{1}{2}[H_0, P(K)]_+ = (dV + 3N_1(K))P(K) + \sum_{\langle x, y \rangle} (P(Q_+(x, y)K) - P(Q_-(x, y)K)). \quad (2.12)$$

The sum on the right hand side extends over configurations which are obtained from the configuration $K = [x, y_1][x_2, y]\dots$ by the following permutations:

$$Q_+(x, y)[x, y_1][x_2, y] = [x, y][x_2, y_1], \quad (2.13)$$

$$Q_-(x, y)[x, y_1][x_2, y] = [y, y_1][x_2, x]. \quad (2.14)$$

Eqs.(2.5), (2.6) and (2.12) will provide us with an evolution equation of the Boltzmann factor between valence bond states.

III. PROJECTION OPERATORS IN SPIN S SECTORS

For the construction of states with total spin S , we follow the original work of Hulthén¹:

$$|K_S\rangle = [z_1, z_2, \dots, z_{2S}]_s [x_1, y_1] \dots [x_v, y_v]. \quad (3.1)$$

Here $[z_1, z_2, \dots, z_{2S}]_s$ denotes a totally symmetrized state of spins at sites $z_1, \dots, z_{2S}$. As before the $[x_j, y_j]$ are valence bond (spin 0) states. Therefore we can characterize each of these states by a configuration K_S with $v = V/2 - S$ valence bonds and $2S$ 'monomers'. Each site of the lattice is occupied either by one valence bond or one monomer. The corresponding projection operators

$$P(K_S) = |K_S\rangle\langle K_S|, \quad (3.2)$$

have the following properties:

$$\text{tr } P(K_S) = 1, \quad (3.3)$$

$$\sum_{K_S} P(K_S) = a_S 1_S, \quad (3.4)$$

where 1_S is the projection operator onto the sector with total spin S and:

$$a_S = \frac{\nu_S(V)}{n_S(V)}. \quad (3.5)$$

$\nu_S(V)$ is the number of monomer-valence bond configurations K_S and $n_S(V)$ the number of spin S states on a lattice with V sites. The nearest neighbour Heisenberg Hamiltonian projected onto the spin S sector

$$H_S = 1_S \sum_{\langle x, y \rangle} 4(x, y) 1_S, \quad (3.6)$$

can be expanded in terms of the projection operators $P(K_S)$:

$$H_S = \frac{V+2}{a_S} \sum_{K_S} N_1(K_S) P(K_S), \quad (3.7)$$

where

$$N_1(K_S) = \sum_{\langle x, y \rangle} \delta([x, y] \in K_S) \quad (3.8)$$

is the number of dimers on the configuration K_S . The Hamiltonian H_S obeys anticommutation relations with the projection operators $P(K_S)$, which are proven in appendix A as well:

$$\begin{aligned} \frac{1}{2}[H_S, P(K_S)]_+ &= (dV + 3N_1(K_S) - 4N_2(K_S))P(K_S) \\ &+ \sum_{\langle x, y \rangle} (P(Q_+(x, y)K_S) - P(Q_-(x, y)K_S)). \end{aligned} \quad (3.9)$$

$N_2(K_S)$ is the number of monomer pairs on neighbouring sites $\langle x, y \rangle$. The action of the permutation operators $Q_{\pm\sigma}(x, y)$ on the valence bond states is given in eqs.(2.13-14) and depicted in Fig.1a). If monomers are involved, one has the following rules:

$$Q_+(x, z)[z, z_2, \dots]_s [x, y] = [y, z_2, \dots]_s [x, z] \quad (3.10)$$

$$Q_-(x, z)[z, z_2, \dots]_s [x, y] = [x, z_2, \dots]_s [z, y], \quad (3.11)$$

which are visualized in Fig.1b). The sum on the right hand side of eq. (3.9) does not include nearest neighbours $\langle x, y \rangle$, which are occupied by dimers or monomer pairs. Finally there exists the analogue of eq.(2.8) for the expansion of the spin-spin correlators projected on the spin S sector:

$$1_S(x, y) 1_S = \frac{V+2}{4a_S} \sum_{K_S} \delta([x, y] \in K_S) P(K_S). \quad (3.12)$$

IV. EXPECTATION VALUES OF THE BOLTZMANN FACTOR BETWEEN MONOMER-VALENCE BOND STATES

Let us introduce the following notation for the eigenvalues h_S and eigenstates $|h_S\rangle$ of H_S :

$$H_S|h_S\rangle = h_S V|h_S\rangle. \quad (4.1)$$

Furthermore let us consider the expectation values of the Boltzmann factor $\exp(\beta H_S)$ between monomer-valence bond states $|K_S\rangle$:

$$f(\beta, K_S) = \langle K_S | \exp(\beta H_S) | K_S \rangle = \sum_{h_S} \exp(\beta V h_S) \langle h_S | P(K_S) | h_S \rangle. \quad (4.2)$$

Then eq.(3.9) -evaluated between the eigenstates $|h_S\rangle$ - yields a system of linear differential equations describing the ‘evolution’ of the expectation values $f(\beta, K_S)$:

$$\begin{aligned} \frac{\partial}{\partial \beta} f(\beta, K_S) &= (dV + 3N_1(K_S) - N_2(K_S)) f(\beta, K_S) \\ &+ \sum_{\langle x, y \rangle} (f(\beta, Q_+(x, y) K_S) - f(\beta, Q_-(x, y) K_S)), \end{aligned} \quad (4.3)$$

with the initial condition:

$$f(\beta = 0, K_S) = 1, \quad \text{for all } K_S. \quad (4.4)$$

The expectation values $f(\beta, K_S)$ contain the whole information on the thermodynamical and magnetic properties of the system:

$$\text{tr} (\exp \beta H_S) = \frac{1}{a_S} \sum_{K_S} f(\beta, K_S). \quad (4.5)$$

Eq.(4.5) results from the ‘completeness’ relation (3.4) for the projection operators $P(K_S)$. Similarly one gets for the spin-spin correlators (x, y) , weighted with the Boltzmann factor:

$$\text{tr} [(x, y) \exp \beta H_S] = \frac{V+2}{4a_S} \sum_{K_S} \delta([x, y] \in K_S) f(\beta, K_S) \quad (4.6)$$

as can be derived from the representation (3.12). In the combined limit:

$$V, S \rightarrow \infty, \quad M = S/V, \quad \beta \text{ fixed} \quad (4.7)$$

the free energy per site at fixed inverse temperature β and magnetization M is given by:

$$F(\beta, M) = \frac{1}{\beta V} \log \text{tr} (\exp \beta H_S). \quad (4.8)$$

Note that in the absence of an external magnetic field the magnetization M (in the antiferromagnetic system) is zero. This means that the free energy $F(\beta, M = \frac{S}{V} = 0)$ in the spin $S = 0$ sector contains already the whole information on the thermodynamics in the absence of an external field. Of course the same holds for the free energies in any of the sectors with fixed total spin S and for consistency they have to be the same. This can be seen immediately from the evolution equations (4.3), which differ for the various values of S by the number of monomers in the configurations K_S . It is at least plausible that a finite number of monomers will not change the behavior of the $f(\beta, K_S)$ in the thermodynamical limit.

V. THE MONOMER DIMER CLUSTER EXPANSION

In this section, we want to investigate an approximation scheme for the evolution equation (4.3). Let us first ask, what are the most important characteristics of the monomer valence bond configuration K_S which determine the variation of $f(\beta, K_S)$ with K_S . Obviously, such characteristic variables are the densities of dimers and monomer pairs $x_1 = N_1/V$, $x_2 = N_2/V$ which appear explicitly in the evolution equation (4.3). It is clear from eq.(4.3) that the $f(\beta, K_S)$ increase much stronger with β on configurations with high dimer- and low monomer pair density. Next the clustering of the monomers and dimers seems to be important as will be demonstrated below. Therefore, the expectation values $f(\beta, K_S)$ are supposed to depend on the numbers $N_1(K_S)$, $N_2(K_S)$, ... of dimers, monomer pairs, neighbouring dimer pairs, etc.

Let us discuss first the lowest order in this approximative scheme, where we average the evolution equation (4.3) over all monomer valence bond configurations with fixed numbers $N_1(K_S)$, $N_2(K_S)$ of dimers and monomer pairs. The approximation would be exact if the right hand side could be expressed as well in terms of the averages

$$f(\beta, N_1, N_2) = \frac{1}{\nu(N_1, N_2, V)} \sum_{K_S(N_1, N_2)} f(\beta, K_S(N_1, N_2)). \quad (5.1)$$

This is not possible for all the terms in the sum on the right hand side of eq.(4.3). Here we approximate the $f(\beta, K_S)$ by their average (5.1). The approximate evolution equation can be brought in the form:

$$\begin{aligned} \frac{\partial}{\partial \beta} f(\beta, N_1, N_2) &= (dV + 3N_1 - N_2) f(\beta, N_1, N_2) \\ &+ V \sum_{j_1, j_2} c_{j_1, j_2}(N_1, N_2, V) f(\beta, N_1 + j_1, N_2 + j_2). \end{aligned} \quad (5.2)$$

The coefficients $c_{j_1, j_2}(N_1, N_2, V)$ are given by :

$$\begin{aligned} c_{j_1, j_2}(N_1, N_2, V) &= \frac{1}{V \nu(N_1, N_2, V)} \sum_{\langle x, y \rangle} \sum_{K(N_1, N_2)} \sum_{\sigma=+, -} \\ &\sigma \prod_{l=1, 2} \delta(N_l(K) + j_l - N_l(Q_\sigma(x, y)K)). \end{aligned} \quad (5.3)$$

Note that the permutations $Q_\pm(x, y)$ (eqs.(2.13,14),(3.10,11)) -applied on the monomer valence bond configuration $K_S(N_1)$ - can change the number of dimers and monomer pairs:

$$N_1(Q_\pm(x, y)K_S) - N_1(K_S) = j_{1\pm}(K_S) = -2, -1, 0, 1, 2 \quad (5.4)$$

$$N_2(Q_\pm(x, y)K_S) - N_2(K_S) = j_{2\pm}(K_S) = -(d-1), \dots, (d-1). \quad (5.5)$$

The coefficients (5.3) are investigated in appendix B. They can be expressed in terms of probabilities to find certain clusters on the monomer valence bond configurations $K_S(N_1, N_2)$. In the combined limit:

$$V, S, N_1, N_2 \rightarrow \infty, \quad x_1 = N_1/V, \quad x_2 = N_2/V, \quad \beta, \quad M = S/V \quad \text{fixed}, \quad (5.6)$$

they only depend on the densities x_1, x_2 of dimers and monomerpairs:

$$c_{j_1 j_2}(N_1, N_2, V) \rightarrow c_{j_1 j_2}(x_1, x_2). \quad (5.7)$$

Note, that the magnetization $M = S/V$ - i.e. the monomer densities - is held fixed. It is a conserved quantity in the evolution process.

The coefficients (5.3) acquire a particularly simple form in the limit:

$$M \rightarrow 0, \quad x_2 \rightarrow 0, \quad c_{j_1 j_2}(x_1, x_2) \rightarrow c_{j_1}(x_1) \delta_{j_2 0}, \quad (5.8)$$

where

$$c_{-2}(x_1) = -w_1(x_1), \quad (5.9)$$

$$c_{-1}(x_1) = 3w_1(x_1) - w_2(x_1) - 2(2d-1)x_1, \quad (5.10)$$

$$c_0(x_1) = -3w_1(x_1) + w_2(x_1) + 4(2d-1)x_1 - d + x_1, \quad (5.11)$$

$$c_1(x_1) = w_1(x_1) + d - x_1(2(2d-1) + 1). \quad (5.12)$$

Here $w_1(x), w_2(x)$ are the probabilities to find on the valence bond configuration K_S :

- dimer pairs with arbitrary orientation: $w_1(x_1)$
- dimer pairs with parallel orientation: $w_2(x_1)$.

These types of clusters are shown in Fig. 2a-b for the case $d = 2$.

VI. THE EVOLUTION EQUATION IN THE THERMODYNAMICAL LIMIT

In this section we are going to study the approximate evolution equation (5.2) in the combined limit (5.6). Then (5.2) reduces to a partial differential equation for if we assume that the averages (5.1) have the form:

$$f(\beta, N_1, N_2) = \exp(V\phi(\beta, x_1, x_2)), \quad (6.1)$$

$$\frac{\partial}{\partial x_l} \phi(\beta, x_1, x_2) = \log R_l \quad l = 1, 2, \quad (6.2)$$

$$\frac{\partial}{\partial \beta} \phi(\beta, x_1, x_2) = L(x_1, R_1, x_2, R_2), \quad (6.3)$$

where

$$L(x_1, R_1, x_2, R_2) = d + 3x_1 - x_2 + \sum_{j_1, j_2} c_{j_1, j_2}(x_1, x_2) R_1^{j_1} R_2^{j_2}. \quad (6.4)$$

The initial condition:

$$\phi(\beta = 0, x_1, x_2) = 0 \quad (6.5)$$

is a consequence of eq.(4.4). To solve eqs.(6.2-5) we first introduce -via a Legendre transform- a function:

$$\Phi(x_1, R_1, x_2, R_2) = \phi(\beta, x_1, x_2) - \beta \frac{\partial \phi(\beta, x_1, x_2)}{\partial \beta}, \quad (6.6)$$

which depends on the four variables x_1, x_2, R_1, R_2 . The partial derivatives of Φ with respect to these variables are not independent but obey the following four relations:

$$\beta = - \frac{\partial \Phi}{\partial R_l} \left(\frac{\partial L}{\partial R_l} \right)^{-1} \quad l = 1, 2, \quad (6.7)$$

$$\beta = - \left(\frac{\partial \Phi}{\partial x_l} - \log R_l \right) \left(\frac{\partial L}{\partial x_l} \right)^{-1} \quad l = 1, 2. \quad (6.8)$$

A solution of these equations is found with the ansatz:

$$\begin{aligned} \Phi(x_1, R_1, x_2, R_2) = & x_1 \log R_1 + x_2 \log R_2 - \int_1^{R_1} \int_1^{R_2} \frac{dR'_1 dR'_2}{R'_1 R'_2} g(R'_1, R'_2, L) \\ & - \int_1^{R_1} \frac{dR'_1}{R'_1} g_1(R'_1, L) - \int_1^{R_2} \frac{dR'_2}{R'_2} g_2(R'_2, L), \end{aligned} \quad (6.9)$$

if the integrands $g(R'_1, R'_2, L), g_1(R'_1, L), g_2(R'_2, L)$ with $L = L(x_1, R_1, x_2, R_2)$ satisfy the conditions:

$$x_1 = x_1(R_1, R_2, L) = \int_1^{R_2} \frac{dR'_2}{R'_2} g(R_1, R'_2, L) + g_1(R_1, L), \quad (6.10)$$

$$x_2 = x_2(R_1, R_2, L) = \int_1^{R_1} \frac{dR'_1}{R'_1} g(R'_1, R_2, L) + g_2(R_2, L). \quad (6.11)$$

Note that the ansatz (6.9) satisfies the initial condition (6.5). According to (6.2), the initial point $\beta = 0$ is mapped on the point $R_1 = R_2 = 1$ in the R_1, R_2 -plane. There the Legendre transform (6.6) has to vanish for all x_1, x_2 . Eqs.(6.10,11) define x_1, x_2 as functions of R_1, R_2, L with partial derivatives:

$$R_2 \left. \frac{\partial x_1}{\partial R_2} \right|_{R_1, L} = R_1 \left. \frac{\partial x_2}{\partial R_1} \right|_{R_2, L} = g(R_1, R_2, L). \quad (6.12)$$

Therefore $x_l, l = 1, 2$ can be written as the gradient:

$$x_l = \frac{\partial}{\partial \log R_l} G(R_1, R_2, L) \quad (6.13)$$

of a ‘potential’ $G(R_1, R_2, L)$. Inserting (6.13) into (6.4) we end up with a partial differential equation for $G(R_1, R_2, L)$. The solution of this equation demands the knowledge of the coefficients $c_{j_1, j_2}(x_1, x_2)$ in (6.4). They will be computed in the subsequent papers, where we treat the thermodynamical and magnetic properties of the antiferromagnetic Heisenberg model in dimensions $d = 1, 2, 3$.

VII. SUMMARY AND PERSPECTIVES

The main result of this paper is the evolution equation (4.3) for the expectation values (4.2) of the Boltzmann factor between monomer valence bond states. Once the evolution equation is solved, thermodynamical and magnetic properties (cf.(4.5)) as well as spin-spin correlators (cf.(4.6)) can be easily computed. The high temperature expansion of eq.(4.3) yields a recursion relation in a vector space of dimension $\nu_S(V)$. This is just the number of monomer valence bond configurations. This number can be reduced by using the symmetries of the Hamiltonian H_S . It seems to be realistic that a direct solution of such a recursion relation is feasible on clusters up to 18 sites.

On the other hand, the exact evolution equation (4.3) can be considered as the starting point of a new type of approximation for the quantum Heisenberg model. In the thermodynamical limit the problem of solving the approximated equation (5.2) can be reduced to the solution of a partial differential equation. The approximation of the evolution equation (4.3) can be systematically improved. The prize one has to pay is that the corresponding partial differential equations become more complex.

We meet the most simple situation in the spin 0 sector, where the approximated evolution equation (5.2) can be solved analytically- up to integrations. We will discuss this solution in a second paper and derive from it explicit expressions for the specific heat of the antiferromagnetic Heisenberg model in dimensions $d = 1, 2, 3$.

* e-mail: muetter@wpts0.physik.uni-wuppertal.de

¹ L. Hulthén, Ark. Mat. Astron. Fys. A26 (11), 1 (1938)

² H.J. Schulz and T.A. Ziman, Europhys. Lett. 18, 355 (1992); H.Q. Lin, Phys. Rev. B 42, 6561 (1990); E. Dagotto and A. Moreo, Phys. Rev. B38, 5087 (1989); J. Oitma and D.D. Betts, Can. J. Phys. 56, 897 (1978);

³ Hong-Qiang Ding and Miloje s. Makivic, Phys. Rev. Lett. 64, 1449 (1990); Phys. Rev. B43, 3562 (1991); G. Gomez-Santos, J.D. Joannopoulos and J.W. Negele, Phys. Rev. B39, 4435 (1989); J.R. Reger and A.P. Young, Phys. Rev. B37, 5978 (1988); E. Mansoukis and R. Salvador, Phys. Rev. Lett. 60, 840 (1988); Phys. Rev. B39, 575 (1987)

⁴ U.J. Wiese and H.P. Ying, University of Bern preprint, Feb. (1993); O. Haan, J. U. Klaetke and K.-H. Mütter, Phys. Rev. B46, 5723 (1992)

⁵ J. Borysowicz, P. Horsch and T. Kaplan, Phys. Rev. B31, 1590 (1985); K. Kubo, T. Kaplan and J. Borysowicz, Phys. Rev. B38, 11550 (1988); H.Q. Lin and D.K. Campbell, J. Appl. Phys. 69 (8), 5947 (1991); S. Liang, Phys. Rev. Lett. 64, 1597 (1990); M. Karbach and K.-H. Mütter, Z. Phys. B90, 83 (1993)

⁶ E. Lieb and D.J. Mattis, Math. Phys. 3, 749 (1962)

⁷ P.W. Anderson, B. Doucot and S. Liang, Phys. Rev. Lett. 61, 365 (1988)

⁸ R. Saito, Solid State Commun. 72, 517 (1989)

APPENDIX A: THE PROPERTIES OF THE VALENCE BOND PROJECTION OPERATORS (VBPO)

The properties of the VBPO's listed in eqs.(2.6), (2.8), (2.10) and (2.12) are based on the following anticommutation relations for the projection operators (x, y) defined in eq.(2.1):

$$(x, y)^2 = (x, y) \quad (\text{A1})$$

$$\frac{1}{2}[(1, 2), (2, 3)]_+ = \frac{1}{4}((1, 2) + (2, 3) - (1, 3)) \quad (\text{A2})$$

$$\frac{1}{2}[(1, 2)(3, 4), (2, 3)]_+ = \frac{1}{4}((1, 2)(3, 4) + (1, 4)(2, 3) - (1, 3)(2, 4)). \quad (\text{A3})$$

For the proof of the completeness relation (2.6) it is sufficient to show that the sum over all VBPO's:

$$\sum_K P(K) = \sum_j a_j (\vec{S}^2)^j \quad (\text{A4})$$

can be written as a power series in the square of the total spin $\vec{S}$. Eq.(A4) obviously implies eq.(2.6): The right hand side -applied to any spin 0 state- produces the same state. The left hand side -applied to any state with spin $s > 0$ - yields zero. To verify (A4) we start with the sum:

$$\Sigma_j = \sum_{x_1, \dots, y_j} (x_1, y_1)(x_2, y_2) \dots (x_j, y_j) \quad (\text{A5})$$

over the product of j disconnected valence bonds. We are going to show by complete induction that:

$$\Sigma_j = \sum_{l=0}^{l=j} a_{jl} (\vec{S}^2)^l \quad (\text{A6})$$

has the property we are looking for. Eq.(A6) is obviously correct for $j = 1$:

$$\Sigma_1 = \sum_{x \neq y} (x, y) = a_{10} + a_{11} \vec{S}^2. \quad (\text{A7})$$

According to the definition (A5), the anticommutator $[\Sigma_j, \Sigma_1]_+$ contains for types of contributions:

- disconnected: The sites x, y do not overlap with the sites $x_1 \dots y_j$. The sum over these contributions yields Σ_{j+1} .
- one site connected: One of the sites x or y overlaps with the sites $x_1 \dots y_j$. The sum over these contributions leads back to terms of the type Σ_j , as can be verified with the anticommutator (A2).
- two site connected: Both sites x and y overlap with $x_1, \dots, y_j$. Again the sum over these contributions leads back to terms of the type Σ_j as can be seen from the anticommutator (A3).

This completes the proof of eq.(A4).

Next we turn to the proof of the representation (2.8). From eqs.(A1) and (A3) we get anticommutation relations for the VBPO's on two types of configurations:

- K_1 : sites x and y are occupied by the same valence bond

$$\frac{1}{2}[P(K_1), (x, y)]_+ = P(K_1) \quad (\text{A8})$$

- K_2 : sites x and y are occupied by two different valence bonds

$$\frac{1}{2}[P(K_2), (x, y)]_+ = \frac{1}{4}(P(K_2) + P(Q_+(x, y)K_2) - P(Q_-(x, y)K_2)). \quad (\text{A9})$$

The action of the permutations $Q_{\pm}\sigma(x, y)$, is defined in eqs.(2.13,14). From eq.(A9) we get

$$[P(K_2) + P(Q_-(x, y)K_2), (x, y)]_+ = P(Q_+(x, y)K_2). \quad (\text{A10})$$

and by summation over all configurations of the type K_2

$$\sum_K [P(K), (x, y)]_+ - \sum_{K_1} [P(K_1), (x, y)]_+ = \frac{V-2}{2} \sum_{K_1} P(K_1). \quad (\text{A11})$$

On the right hand side we have used the fact that the configuration $Q_+(x, y)K_2$ is of the type K_1 . Evaluation of the left hand side by means of the completeness relation (2.6) and the projector property (A8) leads to the representation (2.8). Finally, the anticommutation relations (2.12) for the Hamiltonian H_0 with the VBPO's are derived from eqs.(A8) and (A9) by summation over all nearest neighbour valence bonds (x, y) .

The properties (3.4), (3.9) and (3.12) of the projection operators $P(K_S)$ can be derived in a similar way. For the proof of the completeness relation (3.4) we express the projection operators:

$$|[z_1, \dots, z_{2S}]_+\rangle \langle [z_1, \dots, z_{2S}]_+| = c \prod_{j=1}^{S-1} (\vec{s}^2 - j(j+1)) \quad (\text{A12})$$

as a power series of the square of the total spin $\vec{s}$:

$$\vec{s} = \sum_{l=1}^{2S} \vec{s}(z_l) \quad (\text{A13})$$

on the sublattice which is occupied by the monomers at sites $z_1, \dots, z_{2S}$. From the decomposition (A6) we know, that any power of $\vec{s}^2$:

$$(\vec{s}^2)^j = \sum_{l=0}^j c_{jl} \Sigma'_l \quad (\text{A14})$$

can be written as a sum over terms Σ'_l of the type (A5). In Σ'_l we sum over the product of l valence bonds on the sublattice with sites $z_1, \dots, z_{2S}$.

The sum over all projection operators $P(K_S)$:

$$\sum_{K_S} P(K_S) = \sum_{j=0}^{\frac{V}{2}} b_j \Sigma_j = \sum_{j=0}^{\frac{V}{2}} c_j (\vec{S}^2)^j \quad (\text{A15})$$

is then expressed in terms of the quantities Σ_j defined on the whole lattice. Eq.(A5) yields the expansion in powers of $\vec{S}^2$, which is sufficient for the proof of the completeness relation (3.4).

For the proof of the anticommutation relation (3.9) and the representation (3.12) we have to generalize eqs.(A8) and (A9) to the corresponding equations for the projection operators $P(K_S)$. Here, we have to consider three types of configurations:

- K_1 : sites x and y are occupied by the same valence bond

$$\frac{1}{2} [P(K_1), (x, y)]_+ = P(K_1). \quad (\text{A16})$$

- K_2 : sites x and y are occupied either by two different valence bonds or by one valence bond and one monomer

$$\frac{1}{2} [P(K_2), (x, y)]_+ = \frac{1}{4} (P(K_2) + P(Q_+(x, y)K_2) - P(Q_-(x, y)K_2)). \quad (\text{A17})$$

The action of the permutations $Q_{\pm}(x, y)$ is defined for the first case in (2.13,14) and for the second one in (3.10,11).

- K_3 : sites x and y are occupied by two monomers

$$\frac{1}{2}[P(K_3), (x, y)]_+ = 0. \quad (\text{A18})$$

From eq.(A17) we get

$$[P(K_2) + P(Q_-(x, y)K_2), (x, y)]_+ = P(Q_+(x, y)K_2), \quad (\text{A19})$$

and by summation over all configurations K_2

$$\begin{aligned} \sum_{K_S} [P(K_S), (x, y)]_+ - \sum_{K_1} [P(K_1), (x, y)]_+ - \sum_{K_3} [P(K_3), (x, y)]_+ \\ = \frac{V-2}{4} \sum_{K_1} P(K_1). \end{aligned} \quad (\text{A20})$$

Finally, the representation (3.12) and the anticommutation relation (3.9) follow from (A16), (A18), (A20) and the completeness relation (3.4).

APPENDIX B: MONOMER DIMER CLUSTERS ON THE CONFIGURATIONS K_S

We are going now to exploit equation (5.3) for the coefficients in the approximated evolution equation (5.2). The first sum on the right hand side of eq.(5.3) extends over those nearest neighbours $\langle x, y \rangle$ which are not occupied by a dimer or a monomer pair. The following situations – which are depicted for $d = 2$ and $\langle x, y \rangle = \langle 2, 3 \rangle$ in Fig. 2a-f)-are left:

- K_a : dimer-dimer (not parallel)
- K_b : dimer-dimer (parallel)
- K_c : dimer-valence bond
- K_d : valence bond - valence bond
- K_e : dimer-monomer
- K_f : valence bond-monomer

The valence bonds in the situations K_c, K_d, K_f must not be dimers in order to avoid double counting. On the configurations K_a, K_b, K_c, K_d the neighbouring sites $\langle x, y \rangle = \langle 2, 3 \rangle$ are occupied only by dimers and valence bonds. Therefore the permutations $Q_{\pm}(2, 3)$ will not move the monomers on the configurations, which means that the number of monomer pairs does not change:

$$N_2(Q_{\pm}(2, 3)K_l) - N_2(K_l) = 0 \quad \text{for } l = a, b, c, d. \quad (\text{B1})$$

The changes of the number of dimers (eq.(5.4)) on these configurations can be read of table I.

TABLE I.

	K_a	K_b	K_c	K_d	$K_e(\mu(1), \mu(2), \mu(3))$	$K_f(\mu(x), \mu(2), \mu(3))$
j_{1+}	-1	0	0	1	0	0
j_{1-}	-2	-2	-1	0	-1	0
j_{2+}	0	0	0	-1	$\mu(1) - \mu(3)$	$\mu(x) - \mu(3)$
j_{2-}	0	0	0	-1	$\mu(2) - \mu(3) - 1$	$\mu(2) - \mu(3) - 1$

Let us now turn to the configurations K_e where monomers are involved. The permutations $Q_+(2, 3)$ and $Q_-(2, 3)$ will move the monomers from site 3 to site 1 and site 2, respectively – according to the rules given in Fig. 1b. The resulting change (5.5) in the number of monomer pairs depends on the monomers surrounding the sites 1, 2, and 3 on the configuration $K_e = K_e(\mu(1), \mu(2), \mu(3))$. If we denote by $\mu(x)$, $x = 1, 2, 3$ the number of monomers on the neighbouring sites of x we find:

$$\begin{aligned} N_2(Q_+(2, 3)K_e) - N_2(K_e) &= \mu(1) - \mu(3), \\ N_2(Q_-(2, 3)K_e) - N_2(K_e) &= \mu(2) - \mu(3) - 1. \end{aligned} \quad (\text{B2})$$

In the same way, one can see that the numbers $\mu(x), \mu(2), \mu(3)$ on the neighbouring sites of $x, 2, 3$ determine the change in the number of monomer pairs on the configuration $K_f = K_f(\mu(x), \mu(2), \mu(3))$.

The changes (5.4, 5.5) in the numbers of dimers and monomer pairs – due to the action of the permutations $Q_+(2, 3)$ and $Q_-(2, 3)$ on the configuration K_l $l = a, \dots, f$ – is summarized in table I.

Next we need the probabilities w_l ($l = a, \dots, f$) to find these situations K_l on the monomer valence bond configurations $K_S(N_1, N_2)$. In principal, they can be determined by a Monte Carlo simulation of the monomer valence bond system. The situation simplifies considerably in the spin 0 sector, where we have no monomers and where the complicated situations K_e and K_f do not occur. The remaining situations $K_a - K_d$ have probabilities:

$$w_a = w_1 - w_2, \quad (\text{B3})$$

$$w_b = w_2, \quad (\text{B4})$$

$$w_c = 2(2d - 1)x_1 - 2w_1, \quad (\text{B5})$$

$$w_d = (d - x_1) - w_c - w_1, \quad (\text{B6})$$

$$(\text{B7})$$

which can be completely expressed in terms of the dimer pair probabilities $w_1(x_1), w_2(x_2)$ defined in section V. The coefficients (5.9-12) then follow from table I:

$$c_{-2} = -w_a - w_b, \quad (\text{B8})$$

$$c_{-1} = w_a - w_c, \quad (\text{B9})$$

$$c_0 = w_b + w_c - w_d, \quad (\text{B10})$$

$$c_1 = w_d. \quad (\text{B11})$$

$$(\text{B12})$$

FIGURE CAPTIONS

1. The action of the permutations $Q_\pm(x, y)$ on a monomer-valence bond configuration K_S :
 - a) both sites x and y are occupied by valence bonds
 - b) site z is occupied by a monomer, site x by a valence bond
2. Different types of monomer-valence bond clusters at the nearest neighbour points 2 and 3.




Fig. 1a)




Fig. 1b)









Fig. 2a-f)